

ORIGINAL ARTICLE

Medicine Science 2023;12(2):383-8

The effect of C-reactive protein, Procalcitonin and Neutrophil/lymphocyte ratio on mortality in patients hospitalized in the intensive care unit

 Kemal Yetis Gulsoy¹,  Semiha Orhan²¹Burdur Public Hospital, Intensive Care Unit, Burdur, Türkiye²Afyonkarahisar Health Sciences University, Intensive Care Unit, Afyonkarahisar, Türkiye

Received 13 November 2022; Accepted 13 February 2023

Available online 22.03.2023 with doi: 10.5455/medscience.2022.07.169

Copyright@Author(s) - Available online at www.medicinescience.org

Content of this journal is licensed under a Creative Commons Attribution-NonCommercial-NonDerivatives 4.0 International License.



Abstract

Effectiveness of using inflammatory markers for prognosis assessment in the intensive care units (ICU) is still not clear. The current study aimed to examine the relationship among procalcitonin, C-reactive protein, and neutrophil/lymphocyte ratio of patients who are getting treatment in the ICU during hospitalization and mortality. The study was carried out with a total of 788 patients who were hospitalized in the ICU longer than a day. All participants were over the age of 18 years old. C-reactive protein, procalcitonin, and neutrophil/lymphocyte ratios were compared between the groups of those who died and those who were discharged from the ICU. 54.6% (n= 430) of 788 patients who were admitted to the study, were male whereas 45.4% (n= 358) of them were female. The median age of the study group was calculated as 79 years (IQR = 19 years) (P<0,001). The univariate Cox regression analysis performed to examine the parameters affecting mortality, the analysis suggested following P-values for the procalcitonin, C-reactive protein, and neutrophil/lymphocyte ratio; <0.001, 0.887, and 0.014 respectively. Conversely, the multivariate Cox regression analysis which is performed to examine the parameters affecting mortality yielded following P-values for the same variables respectively: 0.242, 0.116, and 0.523. Although inflammatory markers (i.e., C-reactive protein, procalcitonin, and neutrophil/lymphocyte ratios) were high in patients who died, their relationship with mortality in the ICU could not be shown.

Keywords: SOFA score, APACHE II score, procalcitonin, neutrophil/lymphocyte ratio, intensive care unit

Introduction

Various factors such as age, underlying cause, mechanical ventilation duration, and severity of the disease may affect the mortality rate in the intensive care units (ICU). Different scoring systems are used to evaluate the severity, mortality and morbidity of the disease in ICUs [1]. Two best known and the most used scoring systems among all are the following: The Acute Physiology and Chronic Health Assessment II (APACHE) and The Sequential Organ Failure Assessment (SOFA) [2].

C-reactive protein (CRP) and procalcitonin (PCT) are biomarkers of high specificity that are used as early markers of infection [3]. CRP is an acute phase protein produced by the liver. The plasma concentration level is found in very small amounts in the serum of healthy individuals. Increases in plasma levels occur with stimulation by interleukin 6 during trauma, tissue damage and inflammation stages [4]. CRP, which can be considered as an important indicator of the inflammation stage, interacts with the c-polysaccharide structure in the cell wall of pneumococci and causes collapse. This process activates the classical complement

CITATION

Gulsoy KY, Orhan S. The effect of C-reactive protein, Procalcitonin and Neutrophil/lymphocyte ratio on mortality in patients hospitalized in the intensive care unit. Med Science. 2023;12(2):383-8.



Corresponding Author: Semiha Orhan, Afyonkarahisar Health Sciences University, Intensive Care Unit, Afyonkarahisar, Türkiye
Email: smhorhan@gmail.com

pathway and is the cause of increased phagocytosis [5]. PCT is the precursor of calcitonin produced in the parafollicular-C cells of the thyroid gland. Increased cytokine response with bacterial toxins in systemic bacterial infections stimulates PCT production from non-thyroid parenchymal tissues (kidney, liver, lung, muscle and adipose tissue), causing it to be released into the blood and detected at high levels in serum. PCT values increase to a much lesser extent in cardiogenic shock compared to the large increases seen in patients with septic shock [6]. Studies with healthy volunteers have shown that PCT stimulation is achieved by intravenous injection of a low amount of bacterial endotoxin. PCT begins to be detected in plasma approximately 2-4 hours after bacterial endotoxin injection and reaches a plateau after 6-12 hours with a rapid rise. The PCT level remains high for 24-48 hours, and after reaching its peak value, it returns to its normal level approximately two days later. PCT provides earlier and better results in determining mortality[7]. CRP, Multiple Organ Dysfunction Score (MODS), and APACHE II have been reported as independent factors in determining mortality in ICUs[8].

Neutrophils can be considered as one of the key cell types for innate immune system since they form the very first line of cellular defense system to fight against infection. Lymphocytes are involved in adaptive immunity. The immune response to various infective conditions has different characteristics: increased neutrophil count and decreased lymphocyte count. Against persistent infections, large amounts of neutrophils which may not be apoptotic, are produced. Contrary to lymphocytes, apoptosis of neutrophils in sepsis is beneficial [9]. Lymphocytes are important for an appropriate inflammatory response. The reduction of lymphocytes as a result of apoptosis, cellular depletion, and downregulation enhances the progression of the deleterious inflammatory response [10]. The relationship between neutrophil and lymphocyte concentration was first identified by Zahorec et al.[11]. They expressed the infection as an indicator of the cellular response of the immune system.

Effectiveness of using inflammatory markers for prognosis assessment in the ICU is still not clear yet. Through the current study, we aimed to investigate the relationship between PCT, CRP, and NLR of patients who are getting treatment in the ICU during hospitalization and mortality.

Material and Methods

We pursued this retrospectively planned study in Burdur public hospital ICU. The study took place between the following dates: 01.01.2020 and 01.04.2022. Ethical committee approval, which was obtained in 2022 from the Clinical Research Ethics Committee of Afyonkarahisar Health Sciences University Faculty of Medicine (decision no: 256, dated: 2022), enabled us to conduct the study. All patients participated in the study were over the age of 18. They all were hospitalized and stayed in the ICU longer than 24 hours. Demographic characteristics, laboratory findings, prognosis and mortality data of the patients hospitalized in the ICU were evaluated. Data was obtained through the patient

file information stored in the hospital automation system. The patients were grouped in two: First group included the patients who died in the ICU whereas the second group contained the patients who were discharged. The mean scores of APACHE II, SOFA, CRP, PCT and NLR, obtained during the ICU admission, of two groups (i.e., patients who died, and patients who were discharged) were compared.

Statistical Analysis

Categorical variables obtained from the patients were presented as percentages and frequencies. The Shapiro Wilk test of normality took place to check determine whether the continuous variables deviate from normal distribution. Normally distributed continuous variables were presented in terms of mean (SD), whereas continuous variables do not follow normal distribution were presented in terms of median and interquartile range (IQR). To compare the categorical variables of those who died in the ICU and those who were discharged we conducted the chi-square test. Additionally, we used Fisher's exact test when it was necessary. We used the independent samples t-test for comparing continuous variables between the groups (i.e., patients who died, and patients who were discharged) where the data in each group is normal distributed. In cases where the data does not meet the normal distribution criteria, we applied the Mann-Whitney U tests. Univariate and multivariate Cox regression analyses were used to determine the parameters that increase the risk of mortality. The parameters to be included in the univariate and multivariate analysis were determined. The predictive value of CRP, PCT, NLR, APACHE II and SOFA scores were examined through Receiver Operating Characteristic (ROC), in short ROC, analysis. Optimal cutoff values were determined by the Youden Index.

Results

Of the 788 patients in the study, 430 (54.6 percent of total participant) were male whereas 358 (45.4%) were female. The median of the study group age was calculated as 79 years (IQR=19 years). During the course of the study, 408 (51.8% percent of total participant) patients in ICU's became exitus, while 380 (48.2 percent of total participant) patients were discharged. Of the patients, 96,8% (n=762) had at least one comorbidity, the most common being chronic heart failure (26,7%, n=211), hypertension (26%, n=205), chronic obstructive pulmonary disease (COPD) (20%, n=162), chronic kidney disease (CKD) (16,4%, n=130) diabetes mellitus (15,8%, n=125), sequelae of cerebrovascular disease (CVD) (15,3%, n=121) (Table 1).

The most common reasons for ICU admission were pneumonia (24.3%, n=192), sepsis (20,4%, n=161), decompensated heart failure (14.7% n=116), acute cerebrovascular disease (11.9%, n=94), acute kidney injury (9,9%, n=78), malnutrition (6.8%, n=54), postoperative complication (6,6%, n=52), and trauma (5.2%, n=41) (Table 1).

While PCT was 0.32 (0.09-1.19) ng/mL in patients who were discharged, it was 1.24(0.42-5.45) ng/mL in those who

died ($P<0.001$). CRP (mg/L) was found to be 106.11(52.07-181.55) in patients who died and 55.52(9.77-126) in those who were discharged ($P<0.001$). While NLR was 11.95 (7.30-20.37) in patients who died, it was 8.81(5.22-14) in those who were discharged ($P<0.001$). The comparison of the groups (i.e.,deceased patients vs discharged patients) in terms of characteristics is presented in Table 2.

Table 1. Reasons for ICU admission and comorbidity

	Non-survival (n= 408)	Survival(n= 380)	p
Principal diagnosis leading to ICU admission			
Malnutrition	24(%44.4)	30(%55.6)	0.164**
Pneumonia	114 (%59.4)	78 (%40.6)	0.001**
Sepsis	103 (%64)	58 (%36)	0.001**
AKI	27 (%34.6)	51 (%65.4)	0.001**
Trauma	7 (%17.1)	34 (%82.9)	<0.001**
PC	16 (%30.8)	36 (%69.2)	0.001**
HF	58 (%50)	58 (%50)	0.678**
Acute CVD	59 (%62.8)	35 (%37.2)	0.023**
Comorbidity			
COPD	94 (%58)	68 (%42)	0.074**
HF	116 (%55)	95 (%45)	0.277**
CVD sequelae	75 (%62)	46 (%38)	0.015**
CRF	77 (%59.2)	53 (%40.8)	0.063**
Hypertension	102 (%49.8)	103 (%50.2)	0.501**
DM	64 (%51.2)	61 (%48.8)	0.888**

AKI: Acute kidney injury , CVD: Cerebrovascular disease, HF: Heart failure, CRF: Chronic renal failure, DM:Diabetes mellitus, COPD: Chronic obstructive pulmonary disease, PC: Postoperative complications, *Mann Whitney U test, ** Fisher's exact test

Table 2. Characteristics of the patient groups

	Non-survival (n= 408)	Survival(n= 380)	p
Age(years),median (IQR)	82 (65-85)	76 (65-85)	<0.001*
Sex Male, n(%)	223 (54.7 %)	207 (54.5 %)	1.000**
APACHE II score, median (IQR)	22 (20-27)	18 (15-21)	<0.001*
SOFA score, median (IQR)	9 (7-10)	5 (4-6)	<0.001*
Inflammatory biomarkers, median (IQR)			
Procalcitonin (ng/mL)	1.24 (0.42- 5.45)	0.32 (0.09-1.19)	<0.001*
C-reactive protein (mg/dL)	106.11 (52.07-181.55)	55.52 (9.77- 126)	<0.001*
Neutrophil/lymphocyte ratios	11.95 (7.30-20.37)	8.81 (5.22 – 14)	<0.001*

*Mann Whitney U test, ** Fisher's exact test
APACHE II: Acute Physiology and Chronic Health Assessment II, SOFA: Sequential Organ Failure Assessment

Age, SOFA and APACHE II scores, PCT, CRP and neutrophil/lymphocyte levels were found to be associated with mortality as per the univariate cox regression analysis in our study, while only age, SOFA and APACHE II scores were determined to be the independent risk factors for mortality as per the multivariate Cox regression analysis. The P-value was $P<0.001$ for the PCT, $P=0.887$ for the CRP, and $P= 0.014$ for the NLR in the univariate analysis of Cox regression which was runned to examine the parameters affecting mortality. Moreover, the following p-values were yielded by the multivariate Cox regression

analysis: $P=0.242$, $P=0.116$, and $P=0.523$, respectively. High APACHE II 95% CI 1.03 (1.01-1.05)($P<0.003$) and high SOFA score 95% CI 1.44 (1.37-1.52)($P<0.001$) were obtained. The results of univariate and multivariate Cox regression analyses, which both were performed to examine the parameters affecting mortality, are presented in Table 3.

Table 3. Univariate and multivariate cox regression analysis for mortality

Variables	Univariate			Multivariate		
	HR	%95 GA	P-value	HR	%95 GA	P-value
Age	1.01	1.00-1.02	<0.001	1.01	1.00-1.02	<0.001
APACHE II score	1.13	1.11-1.15	<0.001	1.03	1.01-1.05	0.003
SOFA score	1.49	1.43-1.55	<0.001	1.44	1.37-1.52	<0.001
Procalcitonin	1.01	1.00-1.01	<0.001	1.00	0.99-1.00	0.242
C-reactive protein	1.00	0.99-1.00	0.887			
Neutrophil/lymphocyte ratios	1.00	1.00-1.00	0.014	0.99	0.98-1.00	0.523

HR, Hazardratio. APACHE II: Acute Physiology and Chronic Health Assessment II, SOFA: Sequential Organ Failure Assessment

In ROC analyzes for CRP, PCT, APACHE and SOFA score, NLR, Area Under The Curve(AUC) values were found to be between 0.881 and 0.615. The parameter with the highest AUC value was found to be the SOFA score.

Table 4. Optimal cut-off values and ROC analysis results for CRP, PCT, APACHE II and SOFA score and NLR

Risk factor	AUC(%95)	Cut off	p value	Sensitivity (%)	Specificity (%)
CRP	0.656 (0.613-0.692)	27.12	0.00	0.85	0.39
N/L rate	0.615 (0.573-0.653)	9.71	0.00	0.62	0.57
Prokalsitonin	0.684 (0.642-0.723)	0.64	0.00	0.65	0.66
APACHE score	0.781 (0.749-0.813)	20.50	0.00	0.68	0.74
SOFA score	0.881 (0.862-0.901)	6.51	0.00	0.80	0.78

APACHE II: Acute Physiology and Chronic Health Assessment II, SOFA: Sequential Organ Failure Assessment, CRP: C-reactive protein, NLR: Neutrophil/lymphocyte ratios




Figure 1. ROC analysis for CRP, PCT, APACHE II and SOFA score, NLR. APACHE II: Acute Physiology and Chronic Health Assessment II, SOFA: Sequential Organ Failure Assessment, CRP: C-reactive protein, NLR: Neutrophil/lymphocyte ratios

Discussion

The mortality rate in ICUs is affected by the level of the ICU, co-morbidities of the patients followed, age and many similar factors. In a study conducted in intensive care patients in England, intensive care mortality was reported as 24-41% [12]. Waheed et al. they evaluated the mortality rates of 4165 intensive care patients and found it to be 26.7% [13]. In their study in Turkey, Kaymak et al. reported the average mortality rate as 46.3% in the public, university and private hospital ICUs, while they reported the rate in public hospitals as 47.2% [14]. In our study, the mortality rate of 788 patients hospitalized in ICU of the public hospital was found to be 51.8%.

The SOFA and APACHE II scores are commonly addressed to predict hospital mortality and evaluate multiple organ functions while determining disease severity [15]. The APACHE II score is a widely used scoring system. When the literature is examined, it is seen that there are studies that have a significant relationship between the mortality rate in the ICU and the APACHE II score [16].

The APACHE II score mean was reported as 17.9 (9.4) in the ICON study [17]. In the ICUs of our country; Kamis et al. found the APACHE II score to be 16.8 ± 4.28 in the living group and 23.0 ± 6.22 in the deceased group. Deceased patients had a significantly higher APACHE II score [18]. In the current study, the APACHE II mean score found to be 20 (17-24) for all the patients, with 22 (20-27) in patients who died and 18 (15-21) in patients who were discharged. The study yielded statistically significant mean score difference between deceased patients and discharged patients ($P < 0.001$).

Huang et al., in their cohort study of the sepsis group patients in the ICU, stated that the SOFA value was 13 (10-15) in the patients who died, 6 (4-9) in the living group, and there was a statistical difference between the two groups [19]. In their study in which they evaluated sepsis patients in the ICU of our country, Balcan et al. reported that the average SOFA score of the deceased was $11.5 (\pm 5.2)$, the sofa score of the survivors was $7 (\pm 2.7)$, and that the mortality increased with the increasing SOFA score [20]. In our study, the SOFA score was 9 (7-10) in the group of patients who died and 5 (4-6) in the group of patients who survived ($p < 0.00$). We think that the reason why SOFA values were lower in both groups compared to the studies mentioned in our study was not only the sepsis group but also the heterogeneity of our patient population. No matter how heterogeneous our groups were, a significant difference was found between deceased and living patients' groups. Zou et al. reported the existence of association between both high APACHE II and SOFA scores and mortality [21].

Juan Jesus Rios-Toro et al. reported that changes in the SOFA and APACHE II scoring system were a more sensitive predictor of mortality compared to CRP and PCT in a study they conducted on patients including the sepsis group in the ICU [22]. In their study,

Izabela Kądziołka et al. reported that APACHE II and SAPS II discriminated better in intensive care, had better calibration and were more effective than SOFA as a mortality predictor. They stated that, contrary to the expected results between these scales, SOFA did not reach sufficient effectiveness [23].

The findings of the current study suggested that high SOFA and APACHE II scores might be seen as an independent risk factors for mortality.

CRP is a prognostic factor in many diseases and its relationship with mortality has been reported. Although it has been stated that CRP is associated with mortality, different results have been obtained in the literature regarding its mortality predictive power. While some studies support that CRP may be a good predictor for mortality, there are also studies reporting contrary results [24-25].

One study reported that CRP was associated with mortality in a single variant cox analysis for 28-day mortality in patients hospitalized in the ICU [25]. Wang et al. reported that CRP is an effective biomarker in predicting mortality in their study in unselected patient groups in the ICU. They stated that CRP is an independent risk factor from APACHE II, but it is more effective to evaluate both at the same time [26]. In another study in which they evaluated patients with sepsis, septic shock and SIRS in the ICU, they stated that CRP was not an effective biomarker in predicting hospital mortality [27]. In the current study, we found that CRP value is not an effective biomarker for mortality.

Huang et al. reported that the predictivity of the CRP value for 28-day mortality in the sepsis group in the ICU was AUC 0.656, its sensitivity was 85%, and the cut-off value was 105.4 mg/dL [19]. In our study, the ROC curve values of CRP were AUC 0.656, the sensitivity was 85%, and the cut-off value was 27.1 mg/dL.

Similar to CRP, the increase in PCT also shows the severity of the disease and can be associated with high mortality [28]. Based on the findings of the study investigated the effect of acute-phase reactants on survival in COVID 19 intensive care patients. Orhan et al. [29], suggested PCT level higher than 0.219 ng/mL as an independent risk factor for mortality ($P = 0.017$). Jensen et al. [31], reported PCT as an independent risk factor in determining mortality in their study in which they examined PCT levels in the ICU, but they could not reach the same result for CRP. Zaccone et al. reported the area under the curve of PCT as 0.69 and its sensitivity as 64% in ROC analysis [31]. In our ROC analysis evaluation of our study, PCT was found to be 0.684 AUC and the sensitivity was 65% similar to other studies.

No relationship between CRP and PCT levels and mortality was found in our study. We think that the reason for the difference in results might be due to the heterogeneity of our patient group.

In the current study, patients' serum CRP and PCT values when

they were admitted to the ICU were found to be significantly different between the groups (i.e., deceased patients vs discharged patients). These findings are similar to findings of previously published studies in the literature[30- 32]. Today, the NLR is accepted as a parameter that shows the negative effects of both high neutrophils reflecting acute inflammation and low lymphocyte levels reflecting physiological stress. It has been reported in the literature that NLR is an independent risk factor for mortality and morbidity in various conditions such as cancer and cardiovascular diseases. It is a useful biomarker not only in the case of a significant disease, but also in the detection of postoperative complications, monitoring of the traumatic process, identifying and predicting infectious and inflammatory conditions NLR values in healthy individuals have been reported to be between 0.78-3.53[33].

Heffernan et al. reported that neutrophil count increased and lymphocyte count decreased in SIRS and trauma patients [34]. In their study, Saliccioli et al. reported that the NLR they detected in the samples taken during ICU admission was associated with mortality, but it was not associated with mortality in their subgroups including the sepsis group[35]. In their prospective study, Akilli et al., reported a high rate of NLR in the group they included patients requiring intensive care admission. Their findings suggested it as an independent risk factor for hospital mortality and 6-month mortality [36].

Hwang et al. reported an association of NLR with 28-day mortality when admitted to intensive care. [37]. Martins et al. reported that NLR was an independent risk factor in mortality in their study in the sepsis group[38]. In our study, although the NLR level was higher in patients who died, its relationship with mortality could not be shown.

In their study evaluating the NLR and hospital mortality in patients with sepsis, Ni et al. reported the area under the ROC curve as 0.62 and the sensitivity as 54% [39].

Aygün et al., in their study with patients in the ischemic stroke group in the ICU; they expressed the NLR ratio, the area under the curve in ROC analysis as 0.57, and the sensitivity as 46.9%[40]. In our study, our ROC analysis evaluation; The NLR cut off value was 6.21, the NLR was 0.615, the AUC was drawn and the sensitivity was found to be 62%.

In our ROC analysis results, we found that the most valuable parameter was the SOFA score. Afterwards, they were APACHE II, CRP, PCT and NLR, respectively. Due to the heterogeneity of our patient groups, we could not achieve sufficient statistical performance from the biochemical parameters PCT, CRP and NLR. We believe that scoring systems in ICUs are more useful because they cover many diseases.

Limitations

The first limitation was that the study was single-centred and retrospective. The second limitation of the study was that there

was no categorization performed on the patients according to their clinical diagnoses. This may have led to heterogeneity in our results. The third limitation of the study was shortness of follow-up period. It was not sufficient for evaluation of the patients' long-term prognosis. In the future, it would be beneficial to conduct multicenter prospective studies with a longer follow-up period.

Conclusion

Although inflammatory markers such as PCT, CRP, and NLR were high in patients who died, their relationship with mortality in the ICU could not be shown. The APACHE II and SOFA scores, used as mortality indicators in the ICU, have been determined to be independent risk factors for mortality in our study. It would be more appropriate to use the APACHE II and SOFA scores as an indicator of mortality in the ICU.

Conflict of interests

The authors declare that there is no conflict of interest in the study.

Financial Disclosure

The authors declare that they have received no financial support for the study.

Ethical approval

The study was conducted following the Declaration of Helsinki, and patients gave their written consent. Approval for the study was granted by the Clinical Research Ethics Committee of Afyonkarahisar Health Sciences University (decision no:256, dated:2022)

References

1. Strand K, Flaatten H. Severity scoring in the ICU: a review. *Acta Anaesthesiol Scand.* 2008;52:467-78.
2. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Inten Care Med.* 1996;22:707-10.
3. Henriquez-Camacho C, Losa J. Biomarkers for sepsis. *Bio Med Res Int.* 2014;2014:547818.
4. Simon L, Gauvin F, Amre DK, et al. Serum procalcitonin and C-reactive protein levels as markers of bacterial infection: a systematic review and meta-analysis. *Clin Infect Dis.* 2004;39:206-17.
5. Çolak A, Yılmaz C, Toprak B, et al. Procalcitonin and CRP as biomarkers in discrimination of community-acquired pneumonia and exacerbation of COPD. *J Med Biochem.* 2017;36:122-26.
6. Madhura NS, Shashikala N, Shankar M, et al. Procalcitonin levels versus Microbiological profile in central line associated bloodstream infections (CLABSI) of patients on Hemodialysis. *Asian J Med.* 2021;12:34-42.
7. Carrol ED, Thomson AP, Hart CA. Procalcitonin as a marker of sepsis. *Int J Antimicrob Agents.* 2002;20:1-9.
8. Lobo SMA, Lobo FRM, Bota DP, et al. C-reactive protein levels correlate with mortality and organ failure in critically ill patients. *Chest.* 2003;123:2043-9.
9. Djordjevic D, Rondovic G, Surbatovic M, et al. Neutrophil-to-Lymphocyte Ratio, Monocyte-to-Lymphocyte Ratio, Platelet-to-Lymphocyte Ratio, and Mean Platelet Volume- to-Platelet Count Ratio as Biomarkers in Critically Ill and Injured Patients: Which Ratio to Choose to Predict Outcome and Nature of Bacteremia? *Mediators of Inflammation.* 2018;2018:3758068.
10. Menges T, Engel J, Welters I, et al. Changes in blood lymphocyte

- population safter multiple trauma: association with post traumatic complications. *Crit Care Med.* 1999;27:733–40.
11. Zahorec R. Ratio of neutrophil to lymphocyte counts-rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy.* 2001;102:5-14.
 12. Goldhill DR, Sumner A. Outcome of intensive care patients in a group of British intensive care units. *Crit Care Med.* 1998;26:1337-4.
 13. Waheed U, Williams P, Brett S, et al. White cell count and intensive care unit outcome. *Anaesthesia.* 2003;58:180-3.
 14. Kaymak C, Sencan I, Izdes S, et al. Mortality of adult intensive care units in Turkey using the APACHE II and SOFA systems (outcome assessment in Turkish intensive care units) *Arch Med Sci.* 2018;14:510-5.
 15. Richards G, Levy H, Laterre PF, et al. CURB-65, PSI, and APACHE II to assess mortality risk in patients with severe sepsis and community acquired pneumonia in PROWESS. *J Intensive Care Med.* 2011;26:34-40.
 16. Fadaizadeh L, Tamadon R, Saeedfar K, et al. Performance assessment of Acute Physiology and Chronic Health Evaluation II and Simplified Acute Physiology Score II in a referral respiratory intensive care unit in Iran. *Acta Anaesthesiol Taiwan.* 2012;50:59-62.
 17. Vincent JL, Marshall JC, Namendys-Silva SA, et al. Assessment of the World wide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med.* 2014;2:380-6.
 18. Kandaş A, İşçimen R, Kelebek Girgin N, et al. Prognostic value of microalbuminuria in critically ill patients. *Uludağ Üniversitesi Tıp Fakültesi Dergisi.* 2008;34:107-14.
 19. Huang N, Chen J, Wei Y, et al. Multi-marker approach using C-reactive protein, procalcitonin, neutrophil CD64 index for the prognosis of sepsis in intensive care unit: a retrospective cohort study. *BMC Infectious Diseases.* 2022;22:662.
 20. Balcan B, Olgun Ş, Torlak F, et al. Determination of factors affecting mortality of patients with sepsis in a tertiary intensive care unit. *Türk Toraks Derg.* 2015;16:128-32.
 21. Zou X, Li S, Fang M, et al. Acute Physiology and Chronic Health Evaluation II score as a predictor of hospital mortality in patients of coronavirus disease 2019. *Crit Care Med.* 2020;48:8.
 22. Rios-Toro JJ, Gallego de Guzman MDP, Guerrero- Marin M, et al. Prognostic value of variations in serum biomarkers and prognostic scores values between admission and second day in intensive care unit septic patients. *Cureus.* 2021;13: e16472.
 23. Kądziołka I, Świątek R, Borowska K, et al. Validation of APACHE II and SAPS II scales at the intensive care unit along with assessment of SOFA scale at the admission as an isolated risk of death predictor. *Anaesthesiol Intensive Ther.* 2019;51:107–11.
 24. Prieto MF, Kilstein J, Bagilet D, et al. C-reactive protein as a marker of mortality in intensive care unit. *Med Intensiva.* 2008;32:424-30.
 25. Seller-Pérez G, Herrera-Gutiérrez ME, Lebrón-Gallardo M, et al. Serum C-reactive protein as a marker of outcome and infection in critical care patients. *Med Clin (Barc).* 2005;125:761-65.
 26. Wang F, Pan W, Pan S, et al. Usefulness of N-terminal pro-brain natriuretic peptide and C-reactive protein to predict ICU mortality in unselected medical ICU patients: a prospective, observational study. *Crit Care.* 2011;15:R42.
 27. Schuetz P, Müller B, Nusbaumer C, et al. Circulating levels of GH predict mortality and complement prognostic scores in critically ill medical patients. *Eur J Endocrinol.* 2009;160:157-63.
 28. Assicot M, Gendrel D, Carsin H, et al. High serum procalcitonin concentrations in patients with sepsis and infection. *Lancet.* 1993;341:515-8.
 29. Orhan S, Mecit BBT, Kazan ED, et al. The effect of acute phase reactants on the survival of COVID-19 patients in intensive care. *Progress in Nutrition.* 2021;23:e2021268.
 30. Jensen JU, Heslet L, Jensen TH, et al. Procalcitonin increase in early identification of critically ill patients at high risk of mortality. *Crit Care Med.* 2006;34:2596-602.
 31. Zacccone V, Falsetti L, Nitti C, et al. The Prognostic Role of Procalcitonin in Critically Ill Patients Admitted in a Medical Stepdown Unit: A Retrospective Cohort Study. *Scientific Reports.* 2020;10:4531.
 32. Chirapongsathorn S, Bunraksa W, Chaiprasert A, et al. Adding C-reactive protein and procalcitonin to the model of end-stage liver disease score improves mortality prediction in patients with complications of cirrhosis. *J Gastroenterol Hepatol.* 2018;33:726–32.
 33. Ishizuka M, Shimizu T, Kubota K. Neutrophil-to-lymphocyte ratio has a close association with gangrenous appendicitis in patients undergoing appendectomy. *Int Surg.* 2012;97:299–304.
 34. Heffernan DS, Monaghan SF, Thakkar RK, et al. Failure to normalize lymphopenia following trauma is associated with increased mortality, independent of the leukocytosis pattern. *Crit Care.* 2012;16:R12.
 35. Saliccioli JD, Marshall DC, Pimentel MAF, et al. The association between the neutrophil-to lymphocyte ratio and mortality in critical illness: an observational cohorts study. *Crit Care.* 2015; 19:13.
 36. Akilli NB, Yortanlı M, Mutlu H, et al. Prognostic importance of neutrophil lymphocyte ratio in critically ill patients: short and long term outcomes. *Am J Emerg Med.* 2014;32:1476- 80.
 37. Hwang SY, Shin TG, Jo JJ, et al. Neutrophil to lymphocyte ratio as a prognostic marker in critically-ill septic patients. *Am J Emerg Med.* 2017;35:234-9.
 38. Martins EC, da Fe Silveira L, Viegas K, et al. Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: a case-control study. *Rev Bras Ter Intensiva.* 2019;31:63-70.
 39. Ni J, Wang H, Li Y, et al. Neutrophil to lymphocyte ratio (NLR) as a prognostic marker for in-hospital mortality of patients with sepsis: A secondary analysis based on a single-center, retrospective, cohort study. *Medicine.* 2019;98:46.
 40. Aygun H, Sener A. The effect of neutrophil lymphocyte ratio on mortality in patients followed in the intensive care unit with the diagnosis of ischemic stroke from the emergency department. *Med Sci.* 2022;11:1296-9.